

## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



## International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7:

A61K 49/04, 9/50

(43) International Publication Date: 20 April 2000 (20.04.00)

(21) International Application Number:

PCT/US99/23685

(22) International Filing Date:

12 October 1999 (12.10.99)

(30) Priority Data:

60/103,860

12 October 1998 (12.10.98)

US

(71) Applicant (for all designated States except US): MALLINCK-RODT INC. [US/US]; Lawrence L. Limpus, P.O. Box 5840, 675 McDonnell Boulevard, St. Louis, MO 63134 (US).

(72) Inventor; and

(75) Inventor/Applicant (for US only): CANTRELL, Gary, L. [US/US]; 509 Lakewood Drive, Troy, IL 62294 (US).

(74) Agent: LIMPUS, Lawrence, L.; Mallinckrodt Inc., 675 Mc-Donnell Boulevard, P.O. Box 5840, St. Louis, MO 63134 (US). (81) Designated States: AU, CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

## **Published**

With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: NOVEL ULTRASOUND CONTRAST AGENTS

(57) Abstract

New and structurally distinct ultrasound contrast agents that comprise micron size microbubbles encapsulating a gas within a shell made from a blend of bipolar compounds having inter-molecular hydrophobic regions of mixed carbon chain length. The compounds have the following structure: R<sub>1</sub>-X-Z; R<sub>2</sub>-X-Z; and R<sub>3</sub>-X-Z', where R<sub>1</sub>, R<sub>2</sub> and R<sub>3</sub> are hydrophobic groups selected from the group consisting of straight-chained alkyls, alkylethers, alkylthiolethers, alkyldisulfides, polyfluoroalkyls, and polyfluoroalkylethers having a carbon chain length greater than or equal to 16 and less than or equal to 32 and where R<sub>1</sub> is greater than R<sub>2</sub> and R<sub>1</sub> is greater than or equal to R<sub>3</sub>; R<sub>3</sub> has one or more such hydrophobic groups having the same or different lengths; X is a linker connecting the hydrophobic group to the polar head group; Z is a polar head group selected from the group consisting of CO<sub>2</sub>-M<sup>+</sup>, SO<sub>3</sub>-M<sup>+</sup>, SO<sub>4</sub>-M<sup>+</sup>, PO<sub>3</sub>-M<sup>+</sup>, PO<sub>4</sub>-M<sup>+</sup>2, N(R)4<sup>+</sup>, a pyridinium or substituted pyridinium group, and a zwitterionic group; R is selected from the group consisting of -H, -CH<sub>3</sub>, alkyl, cycloalkyl, substituted cycloalkyls containing one or more heteroatoms, and benzyl and can be the same or different; and Z' is a nonionic group.